arXiv:1006.3362v3 [math-ph] 2 Jul 2010

THE DEGENERATE PARAMETRIC OSCILLATOR AND INCE'S EQUATION

RICARDO CORDERO-SOTO AND SERGEI K. SUSLOV

Abstract. We construct Green's function for the quantum degenerate parametric oscillator in terms of standard solutions of Ince's equation in a framework of a general approach to harmonic oscillators. Exact time-dependent wave functions and their connections with dynamical invariants and $SU(1,1)$ group are also discussed.

1. The Degenerate Parametric Amplifier Hamiltonian

We use standard annihilation and creation operators in the coordinate representation [14], [20], [31] given by

$$\widehat{a}=\sqrt{\frac{m}{2\hbar\omega}}\left(\omega x+\frac{ip}{m}\right),\qquad \widehat{a}^{\dagger}=\sqrt{\frac{m}{2\hbar\omega}}\left(\omega x-\frac{ip}{m}\right) \tag{1.1}$$

with $p=-i\hbar\partial/\partial x$ and $\widehat{a}\widehat{a}^{\dagger}-\widehat{a}^{\dagger}\widehat{a}=1$ throughout this Letter. The following Hamiltonian

$$H(t)=\frac{\hbar\omega}{2}\left(\widehat{a}\ \widehat{a}^{\dagger}+\widehat{a}^{\dagger}\ \widehat{a}\right)-\frac{\hbar\lambda}{2}\left(e^{2i\omega t}\ \widehat{a}^{2}+e^{-2i\omega t}\left(\widehat{a}^{\dagger}\right)^{2}\right) \tag{1.2}$$

of the time-dependent Schrödinger equation

$$i\hbar\frac{\partial\psi}{\partial t}=H(t)\,\psi \tag{1.3}$$

describes the process of degenerate parametric amplification in quantum optics. The first term corresponds to the self-energy of the oscillator representing the mode of interest, the second term describes the coupling of the classical pump to that mode (with the phenomenological constant λ), giving rise to the parametric amplification process (the phase of the pump is taken to be zero at $t=0$ for convenience). Takahasi [45] wrote a detailed paper on this subject as far as in 1965 (see also [24], [33], [35], [36], [37], [38], [39], [41], [47] and references therein). The same Hamiltonian had also been considered by Angelow and Trifonov in the mid 1990s in order to describe the light propagation in the nonlinear $Ti:LiNbO_3$ anisotropic waveguide (details can be found in [1] and [2]). Another form of this variable quadratic Hamiltonian is given by

$$\begin{aligned} H &= \frac{1}{2m}\left(1+\frac{\lambda}{\omega}\cos(2\omega t)\right)p^{2}+\frac{m\omega^{2}}{2}\left(1-\frac{\lambda}{\omega}\cos(2\omega t)\right)x^{2} \\ &\quad +\frac{\lambda}{2}\sin(2\omega t)\,(px+xp),\qquad p=\frac{\hbar}{i}\frac{\partial}{\partial x}. \end{aligned} \tag{1.4}$$

We refer to (1.2)–(1.3) and/or (1.4) as the degenerate parametric oscillator [33]. It can be recognized as a special case of so-called generalized harmonic oscillators that had attracted considerable

---

Date: July 6, 2010.

1991 *Mathematics Subject Classification.* Primary 81Q05, 35C05. Secondary 42A38.

Key words and phrases. The time-dependent Schrödinger equation, generalized harmonic oscillators, degenerate parametric amplification, Green's function, propagator, Ince's equation, Ermakov's equation.

attention over the years in view of their great importance to many advanced quantum problems including coherent states and uncertainty relations, Berry's phase, asymptotic and numerical methods, quantization of mechanical systems, Hamiltonian cosmology, charged particle traps and motion in uniform magnetic fields, molecular spectroscopy and polyatomic molecules in varying external fields. Quadratic Hamiltonians have particular applications in quantum electrodynamics because the electromagnetic field can be represented as a set of forced harmonic oscillators. See, for example, [3], [6], [9], [10], [11], [13], [15], [16], [22], [26], [30], [43], [48], [49] and references therein. Nonlinear oscillators play a central role in the theory of Bose–Einstein condensation [8] because the dynamics of gases of cooled atoms in a magnetic trap at very low temperatures can be described by an effective equation for the condensate wave function known as the Gross–Pitaevskii (or nonlinear Schrödinger) equation [17], [18], [19], [34] and [46].

A goal of this Letter is to construct Green's function of the Hamiltonian (1.4) in the coordinate representation in terms of solutions of Ince's equation studied in [25] and [29] (see also references therein). Exact time-dependent wave functions and connections with linear and quadratic dynamical invariants and $SU(1,1)$ group are also briefly discussed.

2. Generalized Harmonic Oscillators

The degenerate parametric oscillator Hamiltonian (1.4) belongs to quantum systems described by the one-dimensional time-dependent Schrödinger equation

$$i\frac{\partial\psi}{\partial t}=H\psi \tag{2.1}$$

with variable quadratic Hamiltonians of the form

$$H=a(t)\,p^{2}+b(t)\,x^{2}+c(t)\,px+d(t)\,xp,\qquad p=-i\frac{\partial}{\partial x}, \tag{2.2}$$

where $a(t)$, $b(t)$, $c(t)$, and $d(t)$ are real-valued functions of time, t, only (see, for example, [4], [5], [6], [7], [9], [11], [12], [13], [23], [28], [42], [48] and [49] for a general approach and known elementary solutions; a case related to Airy functions is discussed in [21] and our note deals with another special case of transcendental solutions).

The corresponding Green function, or Feynman's propagator, can be found as follows [4], [42]:

$$\psi=G(x,y,t)=\frac{1}{\sqrt{2\pi i\mu_{0}(t)}}\;e^{i\left(\alpha_{0}(t)x^{2}+\beta_{0}(t)xy+\gamma_{0}(t)y^{2}\right)}, \tag{2.3}$$

where

$$\alpha_{0}(t)=\frac{1}{4a(t)}\frac{\mu_{0}'(t)}{\mu_{0}(t)}-\frac{c(t)}{2a(t)}, \tag{2.4}$$

$$\beta_{0}(t)=-\frac{\lambda(t)}{\mu_{0}(t)},\qquad \lambda(t)=\exp\left(\int_{0}^{t}\left(c(s)-d(s)\right)\,ds\right), \tag{2.5}$$

$$\gamma_{0}(t)=\frac{1}{2\mu_{1}(0)}\frac{\mu_{1}(t)}{\mu_{0}(t)}+\frac{c(0)}{2a(0)} \tag{2.6}$$

and $\mu_{0}(t)$ and $\mu_{1}(t)$ are two linearly independent solutions of the characteristic equation

$$\mu''-\tau(t)\,\mu'+4\sigma(t)\,\mu=0 \tag{2.7}$$

with

$$\tau(t) = \frac{a'}{a} + 2c - 2d, \qquad \sigma(t) = ab - cd + \frac{c}{2}\left(\frac{a'}{a} - \frac{c'}{c}\right) \tag{2.8}$$

subject to initial conditions

$$\mu_0(0) = 0, \qquad \mu_0'(0) = 2a(0) \neq 0, \tag{2.9}$$

$$\mu_1(0) \neq 0, \qquad \mu_1'(0) = 0. \tag{2.10}$$

In this note, we present the time-dependent coefficient $\gamma_0(t)$ in terms of two standard solutions μ_0 and μ_1 of the characteristic equation (2.7)–(2.8) with initial data (2.9)–(2.10), respectively, instead of a previously used formula in terms of only one solution μ_0 (see, for example, [4], [6] and [43]). Our expression (2.6) can be verified by a direct differentiation:

$$\left(\frac{\mu_1}{\mu_0}\right)' = \frac{\mu_1'\mu_0 - \mu_1\mu_0'}{\mu_0^2} = \frac{W(\mu_0, \mu_1)}{\mu_0^2}, \tag{2.11}$$

where the Wronskian can be found with the help of Abel's theorem as follows

$$W(\mu_0, \mu_1) = \text{constant } \lambda^2(t)\, a(t). \tag{2.12}$$

Then, as required,

$$\frac{d\gamma_0}{dt} + a(t)\, \beta_0^2(t) = 0 \tag{2.13}$$

and the constant term in (2.6) gives a correct asymptotic as $t \to 0^+$ (cf. Refs. [7] and [42]). The Green function (2.3) is an eigenfunction of the linear dynamical invariant of Dodonov, Malkin, Man'ko and Trifonov [9], [10], [26] and [27] (see also Theorem 1 of [43]).

By the superposition principle the solution of the Cauchy initial value problem can be presented in an integral form

$$\psi(x,t) = \int_{-\infty}^{\infty} G(x,y,t)\ \chi(y)\ dy, \qquad \lim_{t\to 0^+} \psi(x,t) = \chi(x) \tag{2.14}$$

for a suitable initial function χ on $\mathbb{R}$ (a rigorous proof is given in [42] and uniqueness is analyzed in [6]; another form of solution is provided by an eigenfunction expansion [43]; generalized coherent states and transition amplitudes are discussed in a forthcoming paper [44]). In the next sections, we apply these general results to the case of the degenerate parametric oscillator (1.4).

3. Green's Function and Ince Equation

For the Hamiltonian (1.4), variable coefficients are

$$a = \frac{\hbar}{2m}\left(1 + \frac{\lambda}{\omega}\cos(2\omega t)\right), \tag{3.1}$$

$$b = \frac{m\omega^2}{2\hbar}\left(1 - \frac{\lambda}{\omega}\cos(2\omega t)\right), \tag{3.2}$$

$$c = d = \frac{\lambda}{2}\sin(2\omega t) \tag{3.3}$$

and general expressions for Green's function (2.3)–(2.6) can be simplified by letting $\lambda(t) \equiv 1$ and $c(0) = 0$. The characteristic equation is given by

$$\mu'' + \frac{2\lambda\omega \sin(2\omega t)}{\omega + \lambda \cos(2\omega t)}\mu' + \frac{\omega\left(\omega^2 - 3\lambda^2\right) - \lambda\left(\omega^2 + \lambda^2\right)\cos(2\omega t)}{\omega + \lambda\cos(2\omega t)}\mu = 0, \tag{3.4}$$

which can be identified as a special case of the Ince equation [25], [29]:

$$(1 + a_0 \cos 2s)\, y''(s) + b_0 \sin 2s\ y'(s) + (c_0 + d_0 \cos 2s)\, y(s) = 0, \tag{3.5}$$

when $s = \omega t$ and parameters are given by

$$a_0 = \frac{\lambda}{\omega}, \quad b_0 = 2a_0 = 2\frac{\lambda}{\omega}, \quad c_0 = 1 - 3\frac{\lambda^2}{\omega^2}, \quad d_0 = -\frac{\lambda}{\omega}\left(1 + \frac{\lambda^2}{\omega^2}\right). \tag{3.6}$$

Traditionally, a special question which arises in the theory of Ince's equation is the problem of the existence of periodic solutions. By Theorem 7.1 on p. 93 of [25], if Ince's equation (3.5) has two linearly independent solutions of period π, then the polynomial

$$P(\xi) = 2a_0\xi^2 - b_0\xi - d_0/2 \tag{3.7}$$

has a zero at one of the points $\xi = 0, \pm1, \pm2, \ldots$. If (3.5) has two linearly independent solutions of period 2π, then

$$Q(\xi) = 2P(\xi - 1/2) \tag{3.8}$$

vanishes for one of the values $\xi = 0, \pm1, \pm2, \ldots$.

In the case of the degenerate parametric oscillator Hamiltonian (3.6), both of these polynomials are strictly positive:

$$P(\xi) = \frac{2\lambda}{\omega}\left(\left(\xi - \frac{1}{2}\right)^2 + \frac{\lambda^2}{4\omega^2}\right) > 0 \tag{3.9}$$

and

$$Q(\xi) = \frac{4\lambda}{\omega}\left((\xi - 1)^2 + \frac{\lambda^2}{4\omega^2}\right) > 0. \tag{3.10}$$

Thus the corresponding Ince's equation (3.4) may not have two periodic solutions of periods π/ω or $2\pi/\omega$ and our standard solutions μ_0 and μ_1 that satisfy initial conditions (2.9)–(2.10) cannot be constructed in terms of simple variants of the Fourier sine and cosine series:

$$\mu_1(t) = \sum_{n=0}^{\infty} A_{2n} \cos(2\omega n t), \qquad \mu_0(t) = \sum_{n=1}^{\infty} B_{2n} \sin(2\omega n t), \tag{3.11}$$

or

$$\mu_1(t) = \sum_{n=0}^{\infty} A_{2n+1} \cos(\omega(2n+1)t), \qquad \mu_0(t) = \sum_{n=0}^{\infty} B_{2n+1} \sin(\omega(2n+1)t), \tag{3.12}$$

when coefficients A_m and B_m satisfy certain three-term recurrence relations (more details can be found in [25]). Therefore, the degenerate parametric oscillator (1.4) motivates a detailed study of non-periodic solutions of Ince's equation.

4. The Dynamical Invariant and Time-Dependent Wave Functions

Exact time-dependent wave functions of generalized harmonic oscillators (2.1)–(2.2) can also be obtained as eigenfunctions of the quadratic invariant in terms Hermite polynomials and certain solutions of Ermakov-type equation (see Ref. [43] for a modern introduction and references therein). The dynamical invariant of the Hamiltonian (1.4) can be derived from a general form presented in [6], [43] as follows:

$$E(t)=\left(\mu\, p+\frac{m}{\hbar}\frac{\lambda\sin(2\omega t)\,\mu-\mu'}{1+(\lambda/\omega)\cos(2\omega t)}\,x\right)^{2}+\frac{C_0}{\mu^2}\,x^2,\quad p=-i\frac{\partial}{\partial x},\tag{4.1}$$

where $C_0>0$ is a constant and function $\mu(t)$ satisfies the nonlinear auxiliary equation:

$$\begin{aligned}\mu''&+\frac{2\lambda\omega\sin(2\omega t)}{\omega+\lambda\cos(2\omega t)}\mu'+\frac{\omega\left(\omega^2-3\lambda^2\right)-\lambda\left(\omega^2+\lambda^2\right)\cos(2\omega t)}{\omega+\lambda\cos(2\omega t)}\mu\\&=C_0\left(\frac{\hbar}{m}\right)^2\frac{\left(1+(\lambda/\omega)\cos(2\omega t)\right)^2}{\mu^3}.\end{aligned}\tag{4.2}$$

A general solution of this nonlinear equation can be found by the following 'law of cosines':

$$\mu^2(t)=Au^2(t)+Bv^2(t)+2Cu(t)\,v(t)\tag{4.3}$$

in terms of two linearly independent solutions u and v of the homogeneous equation. The constant C_0 is related to the Wronskian of two linearly independent solutions u and v :

$$AB-C^2=C_0\frac{(2a)^2}{W^2(u,v)},\quad W(u,v)=uv'-u'v\tag{4.4}$$

(more details can be found in [43]; see also references therein regarding this so-called Pinney's solution). Thus solution of the corresponding initial value problem is given by

$$\mu^2(t)=\left(\frac{\mu'(0)}{2a(0)}\mu_0(t)+\frac{\mu(0)}{\mu_1(0)}\mu_1(t)\right)^2+\frac{C_0}{\mu^2(0)}\mu_0^2(t),\quad \mu(0)\neq 0\tag{4.5}$$

in terms of standard solutions μ_0 and μ_1 corresponding to initial data (2.9)–(2.10) of the homogeneous equation (4.2).

Time-dependent wave functions can be presented in the form

$$\psi_n(x,t)=e^{-i(n+1/2)\varphi(t)}\;\Psi_n(x,t)\tag{4.6}$$

(see, for example, [43], [49] and references therein), where

$$\frac{d\varphi}{dt}=\sqrt{C_0}\frac{\hbar}{m}\frac{1+(\lambda/\omega)\cos(2\omega t)}{\mu^2},\quad \varphi(0)=0\tag{4.7}$$

and orthonormal time-dependent eigenfunctions $\Psi_n(x,t)$ of the quadratic invariant (4.1) are expressed in terms of Hermite polynomials [32]:

$$\begin{aligned}\Psi_n(x,t)&=D_n\exp\left(ix^2\frac{m}{2\hbar}\frac{(\mu'/\mu)-\lambda\sin(2\omega t)}{1+(\lambda/\omega)\cos(2\omega t)}\right)\\&\times e^{-x^2\sqrt{C_0}/\left(2\mu^2\right)}H_n\left(x\frac{C_0^{1/4}}{\mu}\right),\quad |D_n|^2=\frac{C_0^{1/4}}{\sqrt{\pi}2^n n!\mu}.\end{aligned}\tag{4.8}$$

Then

$$i\frac{\partial \psi_n}{\partial t} = H\psi_n, \qquad E\psi_n = 2\sqrt{C_0}\left(n+\frac{1}{2}\right)\psi_n. \tag{4.9}$$

If the orthonormal initial wave function is given by

$$\psi_n(x,0) = \sqrt{\frac{\varepsilon}{\sqrt{\pi}2^n n!}}\ e^{\left(i\delta-\varepsilon^2/2\right)x^2} H_n(\varepsilon x), \tag{4.10}$$

where ε and δ are constants, then initial conditions for the auxiliary equation (4.2) are given by

$$\mu(0) = \frac{C_0^{1/4}}{\varepsilon}, \qquad \mu'(0) = 2C_0^{1/4}\frac{\hbar}{m}\left(1+\frac{\lambda}{\omega}\right)\frac{\delta}{\varepsilon} \tag{4.11}$$

(the integral $C_0 > 0$ can be chosen at reader's convenience). Then integration of Ermakov's equation (4.2), say with the help of Pinney's solution (4.5), determines complete dynamics of the oscillator-type wave function through the explicit formula (4.8). Further relations of these particular solutions of the Schrödinger equation with the Cauchy initial value problem are discussed in [43] (see Theorem 2).

5. The $SU(1,1)$ Symmetry

The Hamiltonian (1.2) can also be rewritten as follows

$$H(t) = \hbar\omega J_0 - \hbar\lambda\left(e^{2i\omega t}J_+ + e^{-2i\omega t}J_-\right), \tag{5.1}$$

where generators of a non-compact $SU(1,1)$ group are given by

$$J_+ = \frac{1}{2}\left(a^\dagger\right)^2, \qquad J_- = \frac{1}{2}a^2, \qquad J_0 = \frac{1}{4}\left(a^\dagger a + a^\dagger a\right). \tag{5.2}$$

Then a use can be made of the group properties of the corresponding discrete positive series $\mathcal{D}_+^j$ for further investigation of the degenerate parametric oscillator. This is a 'standard procedure' — more details can be found in Refs. [23], [26], [28], [32] and [40] and/or elsewhere.

6. A Special Case

An integrable special case $\lambda = \omega = m = \hbar = 1$ of the degenerate parametric oscillator has been recently considered by Meiler, Cordero-Soto and Suslov [28], [7] (this Hamiltonian is a simplest time-dependent quadratic integral of motion for the linear harmonic oscillator [6]). Here, the Ince equation (3.4) simplifies to

$$\mu'' + 2\tan t\ \mu' - 2\mu = 0. \tag{6.1}$$

It has two elementary non-periodic standard solutions:

$$\mu_0 = \cos t\sinh t + \sin t\cosh t, \qquad \mu_0(0) = 0, \quad \mu_0'(0) = 2, \tag{6.2}$$

$$\mu_1 = \cos t\cosh t + \sin t\sinh t, \qquad \mu_1(0) = 1, \quad \mu_1'(0) = 0 \tag{6.3}$$

and Green's function is given in terms of trigonometric and hyperbolic functions as follows

$$\begin{aligned} G(x,y,t) &= \frac{1}{\sqrt{2\pi i\left(\cos t\sinh t + \sin t\cosh t\right)}} \\ &\quad \times \exp\left(\frac{\left(x^2-y^2\right)\sin t\sinh t + 2xy - \left(x^2+y^2\right)\cos t\cosh t}{2i\left(\cos t\sinh t + \sin t\cosh t\right)}\right) \end{aligned} \tag{6.4}$$

as a simple consequence of expressions (2.3)–(2.6). It is worth noting that formula (6.4) has been obtained in [28] by a totally different approach using the $SU(1,1)$-symmetry of n-dimensional oscillator wave functions and properties of the Meixner–Pollaczek polynomials. More details can be found in [28], [7].

7. An Extension

The degenerate parametric amplification with time-dependent amplitude and phase had been considered by Raiford [38]. The corresponding Hamiltonian, without damping and neglecting high-frequency terms, has the form

$$H(t)=\frac{\hbar\omega}{2}\left(\widehat{a}\,\widehat{a}^{\dagger}+\widehat{a}^{\dagger}\,\widehat{a}\right)-\frac{\hbar\lambda(t)}{2}\left(e^{i(2\omega t+\delta(t))}\,\widehat{a}^{2}+e^{-i(2\omega t+\delta(t))}\left(\widehat{a}^{\dagger}\right)^{2}\right). \tag{7.1}$$

In this model, the phenomenological coupling parameter $\lambda(t)$, which describes the strength of the interaction between the quantized signal of frequency ω and the classical pump of frequency 2ω, and the pump phase $\delta(t)$ are in general functions of time as indicated. It includes the special case of the pump and signal being off-resonance by a given amount ϵ, i. e., the pump frequency being $2\omega+\epsilon$, by letting $\delta(t)=\epsilon t$ and $\lambda(t)=\lambda$, a constant. More details, including Heisenberg's equation of motion for the annihilation and creation operators for the signal photons, can be found in the original paper [38].

In the coordinate representation, the Hamiltonian (7.1) is rewritten as

$$\begin{aligned}H=&\frac{1}{2m}\left(1+\frac{\lambda(t)}{\omega}\cos(2\omega t+\delta(t))\right)p^{2}\\&+\frac{m\omega^{2}}{2}\left(1-\frac{\lambda(t)}{\omega}\cos(2\omega t+\delta(t))\right)x^{2}\\&+\frac{\lambda}{2}\sin(2\omega t+\delta(t))(px+xp),\qquad p=\frac{\hbar}{i}\frac{\partial}{\partial x}\end{aligned} \tag{7.2}$$

and Green's function can be found by using the method described in section 2. Further details are left to the reader.

8. A Conclusion

In this Letter, we have constructed Green's function of the degenerate parametric oscillator (1.4) in terms of standard non-periodic solutions of Ince's equation. To the best of our knowledge, this oscillator was introduced for the first time by Takahasi [45] in order to describe the process of degenerate parametric amplification in quantum optics (see also [24], [33], [35], [37], [38], [41], [47] and references therein). The corresponding Hamiltonian had also been considered later by Angelow and Trifonov [2], [1] in order to describe the light propagation in a nonlinear anisotropic waveguide. Our observation motivates further investigation of properties of the degenerate parametric oscillator including a systematic study of corresponding non-periodic solutions of Ince's equation that seems to be missing in mathematical literature. Moreover, the Hamiltonian (7.1), which was considered by Raiford [38] in the case of the degenerate parametric amplification with time-dependent amplitude and phase, requires a generalization of Ince's equation.

Acknowledgments. We thank Professor Carlos Castillo-Chávez, Professor Vladimir I. Man'ko, Professor Svetlana Roudenko and Professor Dimiter A. Trifonov for support, valuable discussions and encouragement. The authors are indebted to Professor Victor V. Dodonov for kindly pointing out papers [45], [38] and [41] on the degenerate parametric amplification to our attention — his valuable comments have essentially helped to improve the paper. This work is done as a part of the summer 2010 program on analysis of Mathematical and Theoretical Biology Institute (MTBI) and Mathematical, Computational and Modeling Sciences Center (MCMSC) at Arizona State University. The MTBI/SUMS Summer Research Program is supported by The National Science Foundation (DMS-0502349), The National Security Agency (dod-h982300710096), The Sloan Foundation and Arizona State University. One of the authors (RCS) is supported by the following National Science Foundation programs: Louis Stokes Alliances for Minority Participation (LSAMP): NSF Cooperative Agreement No. HRD-0602425 (WAESO LSAMP Phase IV); Alliances for Graduate Education and the Professoriate (AGEP): NSF Cooperative Agreement No. HRD-0450137 (MGE@MSA AGEP Phase II).

REFERENCES

[1] A. Angelow, *Light propagation in nonlinear waveguide and classical two-dimensional oscillator*, Physica A **256** (1998) #7, 485–498.

[2] A. Angelow and D. A. Trifonov, *Schrödinger covariance states in anisotropic waveguides*, Preprint ICTP, IC/95/44 Trieste, Italy, 1995, 15 pp.

[3] M. V. Berry, *Classical adiabatic angles and quantum adiabatic phase*, J. Phys. A: Math. Gen. **18** (1985) # 1, 15–27.

[4] R. Cordero-Soto, R. M. Lopez, E. Suazo and S. K. Suslov, *Propagator of a charged particle with a spin in uniform magnetic and perpendicular electric fields*, Lett. Math. Phys. **84** (2008) #2–3, 159–178.

[5] R. Cordero-Soto, E. Suazo and S. K. Suslov, *Models of damped oscillators in quantum mechanics*, Journal of Physical Mathematics, **1** (2009), S090603 (16 pages).

[6] R. Cordero-Soto, E. Suazo and S. K. Suslov, *Quantum integrals of motion for variable quadratic Hamiltonians*, Annals of Physics, article in press (2010), doi:10.1016/j.aop.2010.02.020; see also arXiv:0912.4900v9 [math-ph] 19 Mar 2010.

[7] R. Cordero-Soto and S. K. Suslov, *Time reversal for modified oscillators*, Theoretical and Mathematical Physics **162** (2010) #3, 286–316; see also arXiv:0808.3149v9 [math-ph] 8 Mar 2009.

[8] F. Dalfovo, S. Giorgini, L. P. Pitaevskii and S. Stringari, *Theory of Bose–Einstein condensation in trapped gases*, Rev. Mod. Phys. **71** (1999), 463–512.

[9] V. V. Dodonov, I. A. Malkin and V. I. Man'ko, *Integrals of motion, Green functions, and coherent states of dynamical systems*, Int. J. Theor. Phys. **14** (1975) # 1, 37–54.

[10] V. V. Dodonov and V. I. Man'ko, *Invariants and correlated states of nonstationary quantum systems*, in: *Invariants and the Evolution of Nonstationary Quantum Systems*, Proceedings of Lebedev Physics Institute, vol. 183, pp. 71-181, Nauka, Moscow, 1987 [in Russian]; English translation published by Nova Science, Commack, New York, 1989, pp. 103-261.

[11] R. P. Feynman, *The Principle of Least Action in Quantum Mechanics*, Ph. D. thesis, Princeton University, 1942; reprinted in: *"Feynman's Thesis – A New Approach to Quantum Theory"*, (L. M. Brown, Editor), World Scientific Publishers, Singapore, 2005, pp. 1–69.

[12] R. P. Feynman, *Space-time approach to non-relativistic quantum mechanics*, Rev. Mod. Phys. **20** (1948) # 2, 367–387; reprinted in: *"Feynman's Thesis – A New Approach to Quantum Theory"*, (L. M. Brown, Editor), World Scientific Publishers, Singapore, 2005, pp. 71–112.

[13] R. P. Feynman and A. R. Hibbs, *Quantum Mechanics and Path Integrals*, McGraw–Hill, New York, 1965.

[14] S. Flügge, *Practical Quantum Mechanics*, Springer–Verlag, Berlin, 1999.

[15] D. Gómez Vergel and E. J. S. Villaseñor, *The time-dependent quantum harmonic oscillator revisited: Applications to quantum theory*, Ann. Phys. **324** (2009), 1360–1385.

[16] J. H. Hannay, *Angle variable holonomy in adiabatic excursion of an integrable Hamiltonian*, J. Phys. A: Math. Gen **18** (1985) # 2, 221–230.

[17] Yu. Kagan, E. L. Surkov and G. V. Shlyapnikov, *Evolution of Bose-condensed gas under variations of the confining potential*, Phys. Rev. A **54** (1996) #3, R1753–R1756.

[18] Yu. Kagan, E. L. Surkov and G. V. Shlyapnikov, *Evolution of Bose gas in anisotropic time-dependent traps*, Phys. Rev. A **55** (1997) #1, R18–R21.

[19] Yu. S. Kivshar, T. J. Alexander and S. K. Turitsyn, *Nonlinear modes of a macroscopic quantum oscillator*, Phys. Lett. A **278** (2001) #1, 225–230.

[20] L. D. Landau and E. M. Lifshitz, *Quantum Mechanics: Nonrelativistic Theory*, Pergamon Press, Oxford, 1977.

[21] N. Lanfear and S. K. Suslov, *The time-dependent Schrödinger equation, Riccati equation and Airy functions*, arXiv:0903.3608v5 [math-ph] 22 Apr 2009.

[22] P. G. L. Leach, *Berry's phase and wave functions for time-dependent Hamiltonian systems*, J. Phys. A: Math. Gen **23** (1990), 2695–2699.

[23] C. F. Lo, *Coherent-state propagator of the generalized time-dependent parametric oscillator*, Europhys. Lett. **24** (1993) #5, 319–323.

[24] W. H. Louisell, *Radiation and Noise in Quantum Electronics*, McGraw-Hill, Inc., New York, 1964.

[25] W. Magnus and S. Winkler, *Hill's Equation*, Dover Publications, New York, 1966.

[26] I. A. Malkin and V. I. Man'ko, *Dynamical Symmetries and Coherent States of Quantum System*, Nauka, Moscow, 1979 [in Russian].

[27] I. A. Malkin, V. I. Man'ko and D. A. Trifonov, *Linear adiabatic invariants and coherent states*, J. Math. Phys. **14** (1973) #5, 576–582.

[28] M. Meiler, R. Cordero-Soto, and S. K. Suslov, *Solution of the Cauchy problem for a time-dependent Schrödinger equation*, J. Math. Phys. **49** (2008) #7, 072102: 1–27; see also arXiv: 0711.0559v4 [math-ph] 5 Dec 2007.

[29] R. Mennicken, *On Ince's equation*, Archive for Rational Mechanics and Analysis **29** (1968) #2, 144–160.

[30] S. Menouar, M. Maamache and J. R. Choi, *An alternative approach to exact wave functions for time-dependent coupled oscillator model of charged particle in variable magnetic field*, Annals of Physics, article in press (2010), doi:10.1016/j.aop.2010.04.011.

[31] E. Merzbacher, *Quantum Mechanics*, third edition, John Wiley & Sons, New York, 1998.

[32] A. F. Nikiforov, S. K. Suslov, and V. B. Uvarov, *Classical Orthogonal Polynomials of a Discrete Variable*, Springer–Verlag, Berlin, New York, 1991.

[33] M. Orszag, *Quantum Optics*, Springer–Verlag, Heidelberg, 1994.

[34] V. M. Pérez-García, P. Torres and G. D. Montesinos, *The method of moments for nonlinear Schrödinger equations: theory and applications*, SIAM J. Appl. Math. **67** (2007) #4, 990–1015.

[35] J. Peřina, Z. Hradil, and B. Jurčo, *Quantum Optics and Fundamentals of Physics*, Kluwer, Dordrecht, 1994.

[36] R. R. Puri, *Mathematical Methods of Quantum Optics*, Springer–Verlag, Berlin, Heidelberg, 2001.

[37] M. T. Raiford, *Statistical dynamics of quantum oscillators and parametric amplification in a single mode*, Phys. Rev. A **2** (1970) #4, 1541–1558.

[38] M. T. Raiford, *Degenerate parametric amplification with time-dependent pump amplitude and phase*, Phys. Rev. A **9** (1974) #5, 2060–2069.

[39] Y. R. Shen, *Principles of Quantum Optics*, Wiley, New York, 1984.

[40] Yu. F. Smirnov and K. V. Shitikova, *The method of K harmonics and the shell model*, Soviet Journal of Particles & Nuclei **8** (1977) #4, 344–370.

[41] D. Stoler, *Photon antibunching and possible ways to observe it*, Phys. Rev. Lett. **33** (1974) #23, 1397–1400.

[42] E. Suazo and S. K. Suslov, *Cauchy problem for Schrödinger equation with variable quadratic Hamiltonians*, under preparation.

[43] S. K. Suslov, *Dynamical invariants for variable quadratic Hamiltonians*, Physica Scripta **81** (2010) #5, 055006 (11 pp); see also arXiv:1002.0144v6 [math-ph] 11 Mar 2010.

[44] S. K. Suslov, *Coherent states and transition amplitudes for generalized harmonic oscillators*, under preparation.

[45] H. Takahasi, *Information theory of quantum-mechanical channels*, in: *Advances in Communication Systems. Theory and Applications*, v. 1, ed (A. V. Balakrishnan, Editor), Academic Publishers, New York, pp. 227–310.

[46] C. Trallero-Giner, J. Drake, V. Lopez-Richard, C. Trallero-Herrero and J. L. Birman, *Bose–Eistein condensates: Analytical methods for the Gross–Pitaevskii equation*, Phys. Lett. A **354** (2006), 115–118.

[47] D. F. Walls and G. J. Milburn, *Quantum Optics*, Springer–Verlag, Berlin, Heidelberg, 1994.

[48] K. B. Wolf, *On time-dependent quadratic Hamiltonians*, SIAM J. Appl. Math. **40** (1981) #3, 419–431.

[49] K-H. Yeon, K-K. Lee, Ch-I. Um, T. F. George and L. N. Pandey, *Exact quantum theory of a time-dependent bound Hamiltonian systems*, Phys. Rev. A **48** (1993) # 4, 2716–2720.

Mathematical, Computational and Modeling Sciences Center, Arizona State University, Tempe, AZ 85287–1904, U.S.A.

E-mail address: `ricardojavier81@gmail.com`

Mathematical, Computational and Modeling Sciences Center, Arizona State University, Tempe, AZ 85287–1904, U.S.A.

E-mail address: `sks@asu.edu`

URL: `http://hahn.la.asu.edu/~suslov/index.html`